A SELECTION OF SCIENTIFIC LITERATURE ON CORD BLOOD PLATELET GEL

© MEDITALIA SRL - REV. 01 - 15-10-2018

- Gelmetti A, Greppi N, Guez S, Grassi F, Rebulla P, Tadini G. Cord blood platelet gel for the treatment of inherited epidermolysis bullosa. Transfus Apher Sci. 2018 Jun;57(3):370-373.
- 2. Volpe P, Marcuccio D, Stilo G, Alberti A, Foti G, Volpe A, Princi D, Surace R, Pucci G, Massara M. Efficacy of cord blood platelet gel application for enhancing diabetic foot ulcer healing after lower limb revascularization. Semin Vasc Surg. 2017 Dec;30(4):106-112.
- 3. Piccin A, Rebulla P, Pupella S, Tagnin M, Marano G, Di Pierro AM, Santodirocco M, Di Mauro L, Beqiri L, Kob M, Primerano M, Casini M, Billio A, Eisendle K, Fontanella F. Impressive tissue regeneration of severe oral mucositis post stem cell transplantation using cord blood platelet gel. Transfusion. 2017 Sep;57(9):2220-2224.
- Sindici E, Astesano S, Fazio L, Dragonetti A, Pugliese M, Scully C, Carossa S, Broccoletti R, Arduino PG. Treatment of Oral Lesions in Dystrophic Epidermolysis Bullosa: A Case Series of Cord Blood Platelet Gel and Low-level Laser Therapy. Acta Derm Venereol. 2017 Mar 10;97(3):383-384.
- 5. Longo V, Rebulla P, Pupella S, Zolla L, Rinalducci S. Proteomic characterization of platelet gel releasate from adult peripheral and cord blood. Proteomics Clin Appl. 2016 Aug;10(8):870-82.
- Rebulla P, Pupella S, Santodirocco M, Greppi N, Villanova I, Buzzi M, De Fazio N, Grazzini G; Italian Cord Blood Platelet Gel Study Group (see Appendix 1). Multicentre standardisation of a clinical grade procedure for the preparation of allogeneic platelet concentrates from umbilical cord blood. Blood Transfus. 2016 Jan:14(1):73-9.
- Parazzi V, Lavazza C, Boldrin V, Montelatici E, Pallotti F, Marconi M, Lazzari L. Extensive Characterization of Platelet Gel Releasate From Cord Blood in Regenerative Medicine. Cell Transplant. 2015;24(12):2573-84.
- 8. Rosso L, Parazzi V, Damarco F, Righi I, Santambrogio L, Rebulla P, Gatti S, Ferrero S, Nosotti M, Lazzari L. Pleural tissue repair with cord blood platelet gel. Blood Transfus. 2014 Jan;12 Suppl 1:s235-42.
- Parazzi V, Lazzari L, Rebulla P. Platelet gel from cord blood: a novel tool for tissue engineering. Platelets. 2010;21(7):549-54.



Intended use

Preparation and administration of cord blood platelet gel

Fields of application

Wound healing

DISCLAMER

This medical device is intended for use by qualified health personnel only. The present information does not constitute and does not intend to constitute recommendations for medical treatment. The company assumes no responsibility for improper use of the device and for incorrect application of the instructions contained in the package leaflet.

The use of blood components is the responsibility of the qualified health personnel.


Advantages of the BioNest system for cord blood fractionation

 pre-assembled set of A-B-C bags (reduced workload and sterile connection costs) • easy opening of bag D


(lower risk of contamination, improved administration)

Product Code 4211000000


MADE IN ITALY



CB is aseptically transferred from the original collection bag into bag "A" by sterile connection




Transfer of PRP into bag "B", sealing, separation and storage at 2-6 °C of bag "A", which contains red blood cells




Transfer platelet-poor plasma (PPP) into bag "C" except the volume necessary to ensure a predefined concentration of the platelets concentrated in the bottom of bag "B" (for example: 1 million per microliter)



Bag "A" is centrifuged with bags "B" and "C" at low speed to sediment the red blood cells and to concentrate the white blood cells into the buffy coat and the platelets into the supernatant platelet rich plasma (PRP)



High speed centrifugation of "B-C" bags



Sealing, separation and freezing of "B" and "C" bags at temperatures below -25 °C


Project design by: Giroidea.it - MI - Italy

© MEDITALIA srl - Rev. 01 - 15-10-2018

© MEDITALIA srl - Rev. 01 - 15-10-2018




CORD BLOOD PLATELET GEL preparation and administration






Sterile connection and transfer





Add 1/3 vol. of 10% Ca gluconate







Top: Easy opening: tear along side engraving

Bottom: CBPG extraction





PATENTED IN ITALY - TO BE PATENTED IN ROW

Address: Via della Piana 1 23030 - Lovero (SO) - ITALY



info@meditaliasrl.com





meditaliasrl.com